arXiv:1303.4319v1 [math.AP] 18 Mar 2013

SEMICLASSICAL CONTROL AND L^2 RESTRICTION BOUNDS FOR NEUMANN DATA ALONG HYPERSURFACES

HANS CHRISTIANSON AND JOHN A. TOTH

Abstract. We prove an h-microlocal control estimate on h^δ-scales ($0 \leq \delta < 1$) for eigenfunctions in a microlocal neighbourhood of a hypersurface H in terms of the mass in the restricted coball bundle and the mass concentrated on the glancing set. We use these control estimates to obtain L^2-restriction bounds for the Neumann data along H.

1. Introduction

We consider here the eigenvalue problem on a compact Riemannian manifold (M, g) with or without boundary, with either Dirichlet or Neumann boundary conditions if $\partial M \neq \emptyset$. That is, we consider

$$\begin{cases} -\Delta_g \varphi_j = \lambda_j^2 \varphi_j, \quad \text{on } M, \\ \langle \varphi_j, \varphi_k \rangle = \delta_{jk} \\ \\ B\varphi_j = 0 \ \text{ on } \ \partial M. \end{cases}$$

Here, $\langle f, g \rangle = \int_M f \bar{g} dV$ is the $L^2(M)$ inner product with respect to the induced Riemannian volume form dV, and where B is the boundary operator, either $B\varphi = \varphi|_{\partial M}$ in the Dirichlet case or $B\varphi = \partial_\nu \varphi|_{\partial M}$ in the Neumann case.

We introduce a hypersurface $H \subset M$, which we assume to be orientable, embedded, and separating in the sense that

$$M \backslash H = M_+ \cup M_-$$

where $M_\pm$ are domains with boundary in M. This is not a restrictive assumption since we can arrange that any hypersurface is part of the boundary of a domain.

Our main result here deals with L^2-restriction bounds for the normalized Neumann data $\lambda^{-1} \partial_\nu \varphi_\lambda |_H$.

Theorem 1. *Suppose $H \subset M$ is a smooth, embedded orientable separating hypersurface and assume that $H \cap \partial M = \emptyset$ if $\partial M \neq \emptyset$. Let $\{\varphi_{\lambda_j}\}_{j=1}^\infty$ denote the L^2-orthonormalized Laplace eigenfunctions on M. Then,*

$$\| \lambda_j^{-1} \partial_\nu \varphi_{\lambda_j} \|_{L^2(H)} = \mathcal{O}(\lambda_j^{\frac{1}{8}}).$$

The result in Theorem 1 generalizes a result of Hassell and Tao [HT] for boundary traces of Dirichlet eigenfunctions (where there is $\mathcal{O}(1)$ upper bound) to *arbitrary* interior hypersurfaces. We note that the universal L^2-restriction upper bound in Theorem 1 for the normalized Neumann data $\lambda^{-1} \partial_\nu \varphi_\lambda |_H$ is substantially better than for the corresponding Dirichlet data $\varphi_\lambda |_H$ which, by [BGT], is only $\mathcal{O}(\lambda^{\frac{1}{4}})$ and the latter estimate is sharp. At this point, we do not know whether the bound for Neumann data in Theorem 1 is sharp. All examples that we are aware of in fact yield normalzed restriction bounds that are $\mathcal{O}(1)$, consistent with the Hassell-Tao estimates for the boundary values of Dirichlet eigenfunctions.

Our result in Theorem 1 relies on a 2-microlocal positive-commutator (Rellich) identity (see Proposition 1.1) which controls the L^2 restriction of the normalized eigenfunction Cauchy data along H:

$$CD(\lambda) := (\varphi_\lambda|_H, \lambda^{-1}\partial_\nu \varphi_\lambda)$$

in terms of interior mass on h^δ-neighbourhoods of the hypersurface. This somewhat technical result applies to any sequence of eigenfunctions (quantum ergodic or not).

In the following we let the semiclasscal parameter $h \in \{\lambda_j^{-1}\}_{j=1}^\infty$ and rescale the Laplacian to the semiclassical operator $P(h) = -h^2\Delta_g$. We abuse notation somewhat and write $\varphi_h = \varphi_\lambda$ for the eigenfunction with eigenvalue $\lambda^2 = h^{-2}$. To state the relevant commutator estimate, we introduce various h-pseudodifferential cutoffs suppressing for the moment some of the technical details. Let $\chi \in C_0^\infty(\mathbb{R}; [0,1])$ with $\chi(u) = 1$ for $|u| \leq 1/2$ and $\chi(u) = 0$ for $|u| > 1$, $\chi_- \in C^\infty(\mathbb{R})$ with $\chi_-(u) = 1$ when $u < -1$ and $\chi_+ \in C^\infty(\mathbb{R})$ with $\chi_+(u) = 1$ when $u > 1$. In addition we require that

$$\chi_-(u) + \chi(u) + \chi_+(u) = 1; \; u \in \mathbb{R}.$$

Let $R(x', \xi') = \sigma(-\Delta_H)(x', \xi')$ be the principal symbol of the induced hypersurface Laplacian $\Delta_H : C^\infty(H) \to C^\infty(H)$ and consider the decomposition of T^*H into 3 pieces given by the radial cutoffs $\chi_{in}, \chi_{tan}, \chi_{out} \in C^\infty(\mathbb{R}; [0,1])$ with $\chi_{in}(x', \xi') = \chi_-(R(x', \xi') - 1), \chi_{tan}(x', \xi') = \chi(R(x', \xi') - 1)$ and $\chi_{out}(x', \xi') = \chi_+(R(x', \xi') - 1)$. Clearly, supp $\chi_{in} \subset B^*H$, supp $\chi_{out} \in T^*H - \overline{B^*H}$ and in addition,

$$\chi_{in}(x', \xi') + \chi_{tan}(x', \xi') + \chi_{out}(x', \xi') = 1; \;\; (x', \xi') \in T^*H. \tag{1.1}$$

For any $\delta \in [0,1)$ we also define the rescaled cutoff functions by $(\chi_{in})_{h,\delta}(x, \xi) = \chi_-(h^{-\delta}(R(x', \xi') - 1)), (\chi_{tan})_{h,\delta}(x, \xi) = \chi(h^{-\delta}(R(x', \xi') - 1))$, and $(\chi_{out})_{h,\delta}(x, \xi) = \chi_+(h^{-\delta}(R(x', \xi') - 1))$.

We denote the corresponding h-Weyl pseudodifferential partition of unity by $(\chi_{tan})^w_{h,\delta} = Op_h^w((\chi_{tan})_{h,\delta})$ and similarly for $(\chi_{out})^w_{h,\delta}$ and $(\chi_{in})^w_{h,\delta}$ (see Figure 1). In the following, we denote the canonical restriction map by $\gamma_H : C^\infty(M) \to C^\infty(H)$ and the corresponding eigenfunction restriction by $\varphi_h^H := \gamma_H \varphi_h \in C^\infty(H)$.

The main technical step in the proof of Theorem 1 is the following small-scale semiclassical Rellich identity. In keeping the notation consistent with that established in the literature, we define

$$\varphi_h^H = \varphi_h|_H, \;\; \varphi_h^{H,\nu} = hD_\nu \varphi_h|_H,$$

where ν is the outward pointing unit normal vector and $D = i^{-1}\partial$ as usual.

Proposition 1.1. *Suppose $H \subset M$ is a smooth, embedded orientable separating hypersurface with interior component M_- and assume $H \cap \partial M = \emptyset$ if $\partial M \neq \emptyset$. Let $\{\varphi_h : h \in \{\lambda_j^{-1}\}\}$ denote the L^2-orthonormalized Laplace eigenfunctions on M. Then for any $\delta \in [0,1)$ and $h \in (0, h_0(\delta)]$ sufficiently small, we have*

$$\left\langle (1 + h^2\Delta_H)(\chi_{in,tan})^w_{h,\delta} \varphi_h^H, \varphi_h^H \right\rangle_{L^2(H)} + \left\langle (\chi_{in,tan})^w_{h,\delta} \varphi_h^{H,\nu}, \varphi_h^{H,\nu} \right\rangle_{L^2(H)}$$
$$= \left\langle b^w_{in,tan}(x, hD_x) \varphi_h, \varphi_h \right\rangle_{L^2(M_-)} + \mathcal{O}(h^\infty).$$


x_n
$\{|\xi| = 1\}$
H
x_n
$\{|\xi'| \leq 1\}$
H
$(\chi_{out})^w_{h,\delta}$
$(\chi_{in})^w_{h,\delta}$
$(\chi_{tan})^w_{h,\delta}$

FIGURE 1. The hypersurface H in Fermi normal coordinates with the restricted cosphere bundle. In the second picture we have projected onto the coball bundle of H and sketched the microlocal partition of unity. The cutoff $(\chi_{in})^w_{h,\delta}$ microlocalizes to the interior of the coball, on scale h^δ from the edge, $(\chi_{out})^w_{h,\delta}$ microlocalizes to the exterior on the same scale, and $(\chi_{tan})^w_{h,\delta}$ (in dashed) microlocalizes to the edge of this set, where $|\xi'| \sim 1$ on scale h^δ. This is the "glancing" set, since these directions have little or no normal direction ξ_n in the lifted cosphere bundle.

Here

$$\begin{aligned} b^w_{in,tan}(x, hD_x) &= b^w_{1,in,tan}(x, hD_x) + b^w_{2,in,tan}(x, hD_x) \\ &= \frac{i}{h}\Big[h^2 D^2_{x_n},\ \chi(x_n)\, hD_{x_n}\, \chi^w_{in,out}\big(\frac{R(x', x_n = 0, \xi') - 1}{h^\delta}\big)\Big] \\ &\quad + \frac{i}{h}\Big[R(x', x_n, hD_{x'}),\ \chi(x_n)\, hD_{x_n}\, \chi^w_{in,out}\big(\frac{R(x', 0, \xi') - 1}{h^\delta}\big)\Big] \end{aligned}$$

(1.2)

$$+\mathcal{O}_\delta(h^{1-\delta})_{L^2 \to L^2},$$

with

$$b_{1,in,tan} \in S^0_{\Sigma_H,\delta} \quad \textit{and} \quad b_{2,in,tan} \in S^\delta_{\Sigma_H,\delta}.$$

The operators $(\chi_\cdot)^w_{h,\delta}$ are in standard operator classes when $\delta \in [0, 1/2)$, but for $\delta \in [1/2, 1)$ they belong to a second microlocal class of semiclassical pseudodifferential operators in $Op^w_h(S^0_{\Sigma_H,\delta})$ (see Definition 2.1). The estimates in Theorem 1.1 in the latter case are more subtle. To state the stronger result, we need to introduce an appropriate class of h-pseudodifferential operators $Op^w_{h,\Sigma_H}(S^0)$ second microlocalized along S^*H and prove a Rellich-type bound for these operators. The appropriate 2-microlocal calculus is treated in section 2.2.2. The closely related calculus in [SjZw1, SjZw2] is carried out in the case where $\Sigma \subset T^*M$ is a hypersurface, but because Fermi coordinates separate base variables near H one can carry out an effective 2-microlocalization along the codimension 2 submanifold of T^*M given by

$$\Sigma_H := (S^*H \oplus N^*_\epsilon H) \cap T^*_H M$$

Here, in terms of Fermi coordinates (x_n, x') near H, $T^*_H M = \{(x', x_n = 0; \xi', \xi_n)\}$, $N^*_\epsilon H = \{(x'; x_n = 0, \xi' = 0, \xi_n); |\xi_n| < \epsilon\}$ and $\Sigma_H = \{(x', x_n = 0; \xi', \xi_n); R(x', \xi') = 1, \, |\xi_n| < \epsilon\}$.

Remark: The results here make no dynamical assumptions, but in the special case where (M, g) is ergodic and $\delta = 0$, the identity in Proposition 1.1 gives the quantum ergodic restriction theorem (QER) in [CTZ] for quantum ergodic sequences of interior eigenfunctions. Apart from the proof of Theorem 1, we believe the identity is of independent interest with other applications as well. For instance, it is an interesting problem to determine whether Proposition 1.1 yields a "small-scale" quantum ergodic restriction theorem when $\delta \in (0, 1)$. We return to this question in a work in progress [CTZ2].

Remark: The estimate in Proposition 1.1 is applied with scale $\delta = 1/4$ in the proof of Theorem 1 and so the control estimate for the full range $\delta \in [0, 1)$ is more general than we need here. However, it has other interesting applications (including small-scale QER) and so we present it here in the more general form.

2. Second-microlocalized pseudodifferential operators

We collect and briefly review here the requisite semiclassical analysis. This includes the various 2-microlocal symbol classes and the corresponding pseudodifferential operator calculus. The material here is a special case of more general two-parameter calculus developed in [SjZw1, SjZw2]. Since our interest lies in establishing L^2-restriction bounds for eigenfunctions along a hypersurface $H \subset M$, we need only consider ambient symbols supported in an $\epsilon > 0$ neighbourhood of H, where $\epsilon > 0$ is arbitrary small. Thus, we introduce Fermi coordinates $x = (x', x_n)$ near H with $x = \exp_{x'}(x_n \nu_{x'})$ with $\nu_{x'}$ an exterior unit normal to H. Since by assumption H is orientable, this is well-defined. In Fermi coordinates, $H = \{x_n = 0\}$ and it is convenient to define our symbols in terms of these coordinates. We do so without further comment.

2.1. **Homogeneous and semiclassical symbol classes.**

We collect for future reference a brief review of the standard symbol classes and corresponding pseudodifferential operators used later on (see also [Zw]). The more subtle 2 microlocal semiclassical analysis is treated in 2.2.2.

The standard homogeneous symbol spaces that are relevant here are

$$S^m_{\rho,\delta}(T^*M)$$

$$= \{a(x,\xi) \in C^\infty(T^*M - 0); |\partial_x^\alpha \partial_\xi^\beta a(x,\xi)| = \mathcal{O}_{\alpha,\beta}(\langle \xi \rangle^{m - \rho|\alpha| + \delta|\beta|}), \, \rho > \delta\} \tag{2.1}$$

As for the semiclassical symbols, the relevant symbol classes for our purposes are

(2.2)

$$S^{m,k}_{cl}(T^*M \times (0,h_0]) = \{a(x,\xi;h) \in C^\infty(T^*M \times (0,h_0]);$$
$$a(x,\xi;h) \sim \sum_{j=0}^{\infty} a_{m-j}(x,\xi) h^{-m+j}, \ a_{m-j} \in S^k_{0,0}\},$$

(2.3)
$$S^m_\delta(T^*M \times (0,h_0])$$
(2.4)
$$= \{a(x,\xi;h) \in C^\infty(T^*M \times (0,h_0]; |\partial_x^\alpha \partial_\xi^\beta a| = \mathcal{O}_{\alpha,\beta}(h^{-m} h^{-\delta(|\alpha|+|\beta|)} \langle \xi \rangle^{-\infty})\},$$

with $\delta \in [0,1)$. Since both the eigenfunctions φ_h and their restrictions $u_h = \varphi_h|H$ have compact h-wavefront sets (see for example [Zw] and Section 5 below) we are interested here in only the case where the ξ variables are in a compact set. Consequently, the semiclassical symbol classes S^m_δ are most relevant. In the special case where $\delta = 0$,

$$S^m_0(T^*M \times (0,h_0]) = \{a \in C^\infty(T^*M \times (0,h_0]); |\partial_x^\alpha \partial_\xi^\beta a| = \mathcal{O}_{\alpha,\beta}(h^{-m} \langle \xi \rangle^{-\infty})\}.$$

When the context is clear, we sometimes just write S^m_δ instead of $S^m_\delta(T^*M \times (0,h_0])$. The case where $\delta = 0$ is sometimes denoted by $S^m(1)$ in the literature.

The corresponding h-Weyl pseudodifferential operators have Schwartz kernels that are sums of the local integrals of the form

$$Op_h^w(a)(x,y) = (2\pi h)^{-n} \int_{\mathbb{R}^n} e^{i\langle x-y,\xi\rangle/h} a(\frac{x+y}{2},\xi;h) d\xi. \tag{2.5}$$

We will use $Op_h^w(a)$, a_h^w and $a^w(x,hD_x)$ interchangeably to denote h-Weyl quantizations of $a(x,\xi;h)$ since each has its advantages. It is standard that for $a \in S^{m_1,k_1}_{cl}, b \in S^{m_2,k_2}_{cl}$,

$$a^w(x,hD_x) \circ b^w(x,hD_x) = e^{ih\sigma(D_x,D_\xi,D_y,D_\eta)/2} a(x,\xi) b(y,\eta)|_{y=x,\eta=\xi}$$
$$= c^w(x,hD_x) \in Op_h^w(S^{m_1+m_2,k_1+k_2}_{cl}(T^*M))$$

with $c(x,\xi;h) = a(x,\xi;h) \# b(x,\xi;h)$ and $\sigma(x,\xi,y,\eta) = y\xi - x\eta$. Similarily, for $a \in S^{m_1}_\delta, b \in S^{m_2}_\delta$ with $\delta \in [0,1/2)$,

$$a^w(x,hD_x) \circ b^w(x,hD_x) = c^w(x,hD_x) \in Op_h^w(S^{m_1+m_2}_\delta(T^*M))$$

with $c(x,\xi;h) = a(x,\xi;h) \# b(x,\xi;h)$.

Since eigenfunctions (and their restrictions) have compact h-wavefront, it is the algebra $Op_h(S^*_\delta)$ that is most relevant here. We point out that for $a^w(x,hD_x) \in Op_h(S^0_\delta)$, $0 \leq \delta < 1/2$, with $a(x,\xi;h) \geq 0$, one also has the sharp Gårding inequality $a^w(x,hD_x) \geq -Ch^{1-2\delta}$ (in the L^2 sense) and indeed the sharper Fefferman-Phong inequality

$$a^w(x,hD_x) \geq -Ch^{2-4\delta} \tag{2.6}$$

also holds [Zw].

2.2. **Semiclassical second-microlocal pseudodifferential cutoffs: microlocal decompostion.**

2.2.1. *Fermi normal coordinates near H.* Frow now on, we let $x = (x', x_n)$ be Fermi normal coordinates in a small tubular neighbourhood $H(\epsilon)$ of H defined near a point $x_0 \in H$. In these coordinates we can locally write

$$H(\epsilon) := \{(x', x_n) \in U \times \mathbb{R}, \, |x_n| < \epsilon\}.$$

Here $U \subset \mathbb{R}^{n-1}$ is a coordinate chart containing $x_0 \in H$ and $\epsilon > 0$ is arbitrarily small but for the moment, fixed. We let $\chi \in C_0^\infty(\mathbb{R})$ be a cutoff with $\chi(x) = 0$ for $|x| \geq 1$ and $\chi(x) = 1$ for $|x| \leq 1/2$. Moreover, in terms of the normal coordinates,

$$-h^2 \Delta_g = \frac{1}{g(x)} hD_{x_n} g(x) hD_{x_n} + R(x_n, x', hD_{x'})$$

where R is a second-order h-differential operator along H with coefficients that depend on x_n, and $R(0, x', hD_{x'})$ is the induced tangential semiclassical Laplacian, Δ_H, on H. Consequently, at the level of symbols,

$$\sigma(-h^2\Delta_g)(x,\xi) = |\xi|_g^2 = \xi_n^2 + R(x', x_n, \xi'),$$

where,

$$\sigma(-h^2\Delta_H)(x', \xi') = R(x', 0, \xi').$$

Let $\gamma_H : C^0(M) \to C^0(H)$ be the restriction operator $\gamma_H(f) = f|_H$. The adjoint $\gamma_H^* : C^0(H) \to C^0(M)$ is then given by

$$\gamma_H^*(g) = g \cdot \delta_H.$$

When there is no risk for confusion we write $\varphi_h^H = \gamma_H \varphi_h$ and similarly $\varphi_h^{H,\nu} = -ih\gamma_H \partial_{x_n} \varphi_h$.

To describe the relevant 2-microlocal h pseudodifferential operators, we first describe the geometry. For $0 < \delta < 1$, the second microlocalization along $S^*H \subset T^*M$ is achieved in several steps.

The first step involves cutting off to a small neighbourhood of the hypersurface

$$\Sigma_1 := \{(x,\xi) \in H(\epsilon); x_n = 0\} = T_H^* M,$$

using a 2-microlocal cutoff $a_1(x_n; h)$ in the normal variable x_n.

Next, we make a full 2-microlocal cutoff to scale h^δ around the hypersurface $\Sigma_2 \subset T^*M$ given by

$$\Sigma_2 := \{(x,\xi) \in H(\epsilon); R(x', x_n = 0, \xi') = 1\}$$

where we note that $R(x', 0, \xi')$ involves coordinates (x', ξ') complementary to (x_n, ξ_n) and so, in particular,

$$\{x_n, R(x', 0, \xi')\} = 0.$$

This allows us to simultaneously 2-microlocalize to the codimension 2 submanifold $\Sigma_H := \Sigma_1 \cap \Sigma_2 \subset T^*M$. In some applications, we localize to $|x_n|, |\xi_n| \leq h^{\delta/2}$, while in others we only localize in an h-independent neighbourhood in x_n of the form $|x_n| \leq h^\delta$. However, the symbols are quite explicit, so no confusion should arise.

The relevant h-neighbourhoods of $\Sigma_j; j = 1, 2$ are , $\Sigma_1(h) = \{(x,\xi) \in H(\epsilon), |x_n| \leq h^{\delta/2}\}$, and $\Sigma_2(h) = \{(x,\xi) \in H(\epsilon), |R(x', 0, \xi') - 1| \leq h^\delta\}$. Finally, the eigenfunction mass 2-microlocalization is concentrated in the set $\Sigma_H(h) = \Sigma_1(h) \cap \Sigma_2(h)$.

Clearly, it follows that as a subset of T^*M, the intersection $\Sigma_1 \cap \Sigma_2 \subset T^*M$ is the codimension-two submanifold $\Sigma_H \subset \mathbb{R}_{\nu_H} \cdot S^*H$ given by

$$\Sigma_H = \{(x', x_n = 0; \xi', \xi_n) \in T^*M; R(x', x_n = 0, \xi') = 1, |\xi_n| < \epsilon\}. \tag{2.7}$$

This is a subset of the total space of the normal line bundle over S^*H with $(x',\xi';v) \in \Sigma_H$ if and only if $(x',\xi') \in S^*H$ and $v = a\nu_{x'}$ with $|a| < \epsilon$ and $\nu_{x'}$ the unit exterior normal at $x' \in H$. Clearly, $\Sigma_H \subset T^*M$ is a submanifold of codimension two and is the total space of a trivial line bundle over the glancing set S^*H, where the trivialization is given by Fermi coordinates $(x', x_n) \in H(\epsilon)$ with $x = \exp_{x'}(x_n \nu_{x'})$.

2.2.2. *Semiclassical pseudodifferential operators second microlocalized along Σ_H.* We introduce here the relevant 2-microlocal algebra of h-pseudodifferential operators localized on very small scales $\sim h^{\delta}$ where $0 < \delta < 1$ that will be used in the proof of our control estimate in Proposition 1.1. When $\delta \in (0,1/2)$ the pseudodifferential calculus is well-known [Zw] but for $\delta > 1/2$ the construction is more subtle. The relevant calculus has been developed in very general framework by Sjöstrand and Zworski [SjZw1, SjZw2] to which we refer the reader for further details. Since a rather simple special case of their calculus will suffice for our purposes, we will attempt to keep the argument fairly self-contained.

Definition 2.1. Let $H \subset M$ be a hypersurface. We say that a semiclassical symbol b is 2-microlocalized along Σ_H and write $b \in S^m_{\Sigma_H,\delta}(T^*M \times (0,h_0])$ provided there exists $\chi \in C_0^\infty(\mathbb{R}), a_1(x_n,\xi_n;h) \in S^0_{\delta/2}(T^*H(\epsilon) \times (0,h_0])$ such that

$$b(x,\xi;h) = h^{-m} a_1(x_n,\xi_n;h) \cdot \chi\left(\frac{R(x', x_n = 0, \xi') - 1}{h^{\delta}} \right), \;\; 0 \leq \delta < 1$$

for all $(x,\xi) \in T^*M$.

Remark. We note that in the specfic case where

$$a_1(x_n,\xi_n;h) = \chi(h^{-\delta}x_n) \cdot \chi(\xi_n), \;\; \chi \in C_0^\infty(\mathbb{R}),$$

by rescaling in phase space $(x_n,\xi_n) \mapsto (h^{\delta/2}x_n, h^{-\delta/2}\xi_n)$ produces a symbol $a_1 \in S^0_{\delta/2}$. In other words, since we are always using the Weyl calculus, it is sufficient that $(\partial_{x_n}\partial_{\xi_n})^k a_1 = \mathcal{O}(h^{-k\delta})$ to be in a good calculus.

We will need the following proposition (see also [SjZw1, SjZw2]).

Proposition 2.2. *Given $a^w(x,hD_x) \in Op_h(S^{m_1}_{\Sigma_H,\delta})$ and $b^w(x,hD_x) \in Op_h(S^{m_2}_{\Sigma_H,\delta})$ it follows that*

$$a^w(x,hD_x) \circ b^w(x,hD_x) = c^w(x,hD_x) \in Op_h(S^{m_1+m_2}_{\Sigma_H,\delta})$$

with

$$c(x,\xi;h) = a(x,\xi,h)\# b(x,\xi,h).$$

Proof. Since

$$d_{\xi'}R(x',0,\xi') \geq C\langle \xi' \rangle, \, C > 0$$

near $\{R(x',0,\xi') = 1\}$, as in [SjZw1, SjZw2], the proof hinges on the following real-principal type quantum normal form construction given in the following Lemma.

Lemma 2.3. *Let $(x_0,\xi_0) \in \Sigma_H$ and let $U \subset T^*H$ be a sufficiently small open neighbourhood of (x_0,ξ_0). For U small, there exists $V \subset T^*\mathbb{R}^{n-1}$ open together with a canonical transformation*

$$\kappa_F : (U, (x_0',\xi_0')) \longrightarrow (V;(0,0)); \;\; \kappa_F(x',\xi') = (y',\eta')$$

and corresponding h-Fourier integral operators $F(h) : C_0^\infty(U) \to C_0^\infty(V)$ *such that*

$$(i)\ F(h)^* \circ \chi_h^w \left(\frac{R(x', 0, \xi') - 1}{h^\delta} \right) \circ F(h) =_{U\times V} \chi_h^w \left(\frac{\eta_1'}{h^\delta}; h \right),$$

$$(ii)\ \ F(h)^* \circ F(h) =_{U\times V} Id,$$

with $\chi \in C_0^\infty(\mathbb{R}^{n-1}; (0,0) \times (0, h_0])$. *Here,* $A(h) =_{U\times V} B(h)$ *denotes h-microlocal equivalence on* $U \times V \subset T^*H \times T^*\mathbb{R}^{n-1}$.

Given Lemma 2.3 one reduces the proof of the proposition to operators in normal form. In the conormal variables (x_n, ξ_n) the composition formula is standard since symbols are in the standard $S^0_{\delta/2}$-classes with $\delta/2 < 1/2$. Since symbols in $S^m_{\Sigma_H,\delta}$ are separable, it suffices to assume that $a(x,\xi;h) = \chi_1(h^{-\delta}(R(x',0,\xi') - 1))$ and $b(x,\xi;h) = \chi_2(h^{-\delta}(R(x',0,\xi') - 1))$ with $\chi_j \in C_0^\infty(\mathbb{R}); j = 1,2$. Let $\psi_{U_j} \times \psi_{V_j} \in Op_h(S_0^0); j = 1, ..., N_0(h)$ be an h-microlocal partition of unity subordinate to a covering of the supports of a and b in T^*H by open sets $U_j; j = 1, ..., N_0(h)$ such that on each U_j Lemma 2.3 holds with h-FIO $F(h)$ with $WF_h'(F(h)) \subset U_j \times V_j$. Of course $N_0(h) = \mathcal{O}(h^{-n\delta})$, which will be absorbed into the $\mathcal{O}(h^\infty)$ error terms. From Lemma 2.3,

$$a_h^w \circ b_h^w = \sum_{j=1}^{N_0(h)} \psi_{U_j}^w F(h) \circ [F(h)^* a_h^w F(h)] \circ [F(h)^* b_h^w F(h)] \circ F(h)^* \psi_{V_j}^w + R(h),$$

where $R(h) \in Op_h(S_0^{-\infty})$. For the inner model operators one simply rescales the fiber variables $(y', \eta'/h^\delta) \mapsto (y'/h^{\delta/2}, \eta'/h^{\delta/2})$ and computes the composition in the model normal coordinates. The result is that in each chart

(2.8)

$$\begin{aligned} &[F(h)^* a_h^w F(h)] \circ [F(h)^* b_h^w F(h)] \\ &\qquad = \big(\chi_1(h^{-\delta}\eta_1') \# \chi_2(h^{-\delta}\eta_1')\big)^w + \mathcal{O}(h^\infty)_{L^2\to L^2} \\ &\qquad = (2\pi h)^{-(n-1)} \int_{\mathbb{R}^{n-1}} e^{i\langle x'-y', \eta'\rangle/h} \psi_{V_j}\Big(\frac{x'+y'}{2h^{\delta/2}}, h^{\delta/2}\eta'\Big)(\chi_1\#\chi_2)(h^{-\delta/2}\eta_1')\, d\eta' \qquad (2.9) \\ &\qquad\quad + \mathcal{O}(h^\infty)_{L^2\to L^2}. \end{aligned}$$

The $\#$ product expansion is computed as usual in the $h^{-\delta/2}$ calculus and then rescaled, which is particularly simple for our special choice of operators:

$$\begin{aligned} (\chi_1\#\chi_2)(h^{-\delta}\eta_1') &= \sum_{j=0}^{\infty} h^j (D_x D_\eta - D_y D_\xi)^j [\chi_1(h^{-\delta}\eta_1') \cdot \chi_2(h^{-\delta}\xi_1')]|_{\xi'=\eta'} \\ &= \chi_1(h^{-\delta}\eta_1') \cdot \chi_2(h^{-\delta}\eta_1') + R(y', \eta'; h), \end{aligned}$$

where, $R(y', \eta'; h) \in S_0^{-\infty}(T^*\mathbb{R}^{n-1})$.

□

Remark. We note that L^2 boundedness for $a_h^w \in Op_h(S^0_{\Sigma_H,\delta})$ with $\delta \in [0,1)$ is clear by passing to normal form. Since $F(h)^*F(h) \cong_{U_j\times V_j} Id$,

$$\begin{aligned} &\|a^w(x, hD_x)\|_{L^2\to L^2} \leq \textstyle\sum_{j=1}^{N_0} \|\chi_h^w(h^{-\delta}\eta_1')\psi^w_{V_j,h}(y',\eta')\|_{L^2\to L^2} \\ &= \textstyle\sum_{j=1}^{N_0} \|\chi^w(h^{1-\delta}\eta_1')\psi^w_{V_j}(y', h\eta')\|_{L^2\to L^2} \leq C_n \|\chi^w \psi^w_{V_j}\|_{L^\infty} + \mathcal{O}(h^{1-\delta}). \end{aligned} \qquad (2.10)$$

The last line follows by rescaling $\eta' \mapsto h\eta'$ in the fiber variables and then applying the Calderon-Vaillancourt theorem.

3. Rellich identity for h-pseudodifferential operators 2-microlocalized along $\Sigma = S^*H$: proof of Proposition 1.1

Our main theorem relies on a Rellich identity which holds for $a^w(x, hD_x) \in Op_h^w(S^0_{\Sigma_H,\delta})$.

Proof of Proposition 1.1. We first assume $H = \partial M_-$ where M_- is a smooth open submanifold of M. This assumption is to be able to use Green's formula, but since we are allowed a cutoff to a subset of H, it is purely technical. We use a Rellich type identity to write the integral of a commutator over M_- as a sum of intergrals over the boundary. The argument is partially motivated by Burq's proof of boundary quantum ergodicity (ie. the case $H = \partial M$) in [Bur] and our earlier joint paper [CTZ].

First, we choose $\psi \in \mathcal{C}_c^\infty(\mathbb{R})$, $\psi(s) \equiv 1$ for $|s| \leq 2$, and we put

$$a^w(x, hD_x) := \chi(x_n)\, \psi_h^w(hD_{x_n})\, hD_{x_n}\, (\chi_{in})^w_{h,\delta}(x', hD_{x'}) \in Op_h^w(S^0_{\Sigma_H,\delta}),$$

where $\chi(u) \equiv 1$ for $|u| \leq 1/2$ and $\chi(u) \equiv 0$ for $|u| > 1$ as in the introduction. Then, a straightforward application of Green's formula and using that $(-h^2\Delta_g - 1)\varphi_h = 0$ yields the Rellich formula

$$\tag{3.1} \begin{aligned} &\frac{i}{h}\int_{M_-} [-h^2\Delta_g,\, a^w(x, hD_x)]\, \varphi_h(x)\overline{\varphi_h(x)}dx \\ &\quad = -\, i \int_H h\partial_{x_n}\, a^w(x', x_n, hD_x)\varphi_h|_H \overline{\varphi_h}|_H d\sigma_H \\ &\qquad + i \int_H a^w(x', x_n, hD_x)\, \varphi_h|_H \overline{h\partial_{x_n}\varphi_h}|_H d\sigma_H \\ &\quad = \int_H hD_{x_n}\, a^w(x', x_n, hD_x)\varphi_h|_H \overline{\varphi_h}|_H d\sigma_H \\ &\qquad + \int_H a^w(x', x_n, hD_x)\, \varphi_h|_H \overline{hD_{x_n}\varphi_h}|_H d\sigma_H. \end{aligned}$$

In the following we continue to use the notation $\varphi_h^H := \varphi_h|_H$ and $\varphi_h^{H,\nu} := hD_\nu\varphi_h|_H$. We begin by computing the RHS of (3.1).

Since $\chi(0) = 1$ and $[\psi_h^w(hD_{x_n}), (\chi_{in})^w_{h,\delta}(x', hD_{x'})] = 0$, it follows that the second term on the RHS of (3.1) is just

$$\tag{3.2} \Big\langle (\chi_{in})^w_{h,\delta}(x', hD_{x'})\, \psi_h^w(hD_{x_n})\, \varphi_h^{H,\nu}, \varphi_h^{H,\nu} \Big\rangle + \mathcal{O}(h^\infty).$$

As for the first term on RHS of (3.1) we get

$$\int_H hD_{x_n}\Big(\chi(x_n)\,\psi_h^w(hD_{x_n})\,hD_{x_n}\,(\chi_{in})^w_{h,\delta}(x',hD_{x'})\varphi_h\Big)\Big|_{x_n=0}\cdot\overline{\varphi_h}|_{x_n=0}\,d\sigma_H \tag{3.3}$$

$$\begin{aligned} &= \int_H \Big[\chi(x_n)\,\psi_h^w(hD_{x_n})\,(\chi_{in})^w_{h,\delta}(x',hD_{x'})\,(hD_{x_n})^2\varphi_h \\ &\quad - ih\chi'(x_n)\,\psi_h^w(hD_{x_n})\,(\chi_{in})^w_{h,\delta}(x',hD_{x'})\,hD_{x_n}\varphi_h\Big]\Big|_{x_n=0}\cdot\overline{\varphi_h}|_{x_n=0}d\sigma_H \end{aligned} \tag{3.4}$$

$$= \int_H \Big[a(x,hD_x)(1-R(x_n,x',hD_{x'}))\varphi_h\Big]\Big|_{x_n=0}\cdot\overline{\varphi_h}|_{x_n=0}d\sigma_H,$$

since $\chi'(0)=0$. Moreover, $((hD_{x_n})^2+R(x,hD_{x'}))\varphi_h=\varphi_h$ in Fermi coordinates and one can add an energy cutoff (modulo $O(h^\infty)$-error) above to the region $\{(x',\xi);\xi_n^2+R(x_n=0,x',\xi')=1+\mathcal{O}(h^\delta)$. But then since $R(x_n=0.x',\xi')=\sigma(-h^2\Delta_H)(x',\xi')\geq 0$, we have $|\xi_n|\gtrapprox 1$ on this set and so the frequency cutoff $\psi_h^w(hD_n)=1$. Consequently, modulo $O(h^\infty)$-error, (3.3) is equal to

$$\int_H \Big[\,(\chi_{in})^w_{h,\delta}(x',hD_{x'})(1-R(0,x',hD_{x'}))\psi_h^w(hD_{x_n})\varphi_h\,\Big)\Big|_{x_n0}\cdot\overline{\varphi_h}|_{x_n=0}d\sigma_H$$

$$=\big\langle(\chi_{in})^w_{h,\delta}(x',hD_{x'})(1+h^2\Delta_H)\varphi_h^H,\varphi_h^H\big\rangle. \tag{3.5}$$

Thus, from (3.2), (3.5) and (3.1), it follows that

$$\begin{aligned} &\Big\langle(\chi_{in})^w_{h,\delta}(x',hD_{x'})\varphi_h^{H,\nu},\varphi_h^{H,\nu}\Big\rangle_{L^2(H)} \\ &\qquad + \big\langle(\chi_{in})^w_{h,\delta}(x',hD_{x'})(1+h^2\Delta_H)\varphi_h^H,\varphi_h^H\big\rangle_{L^2(H)} \\ &= \Big\langle b^w(x,hD_x)\varphi_h,\varphi_h\Big\rangle_{L^2(M_-)}+\mathcal{O}(h^\infty) \end{aligned} \tag{3.6}$$

where,

$$b^w(x,hD_x)=\frac{i}{h}[-h^2\Delta_g,a^w(x,hD_x)]\in Op_h^w(S^\delta_{\Sigma_H,\delta}). \tag{3.7}$$

It follows that

$$\begin{aligned} &b^w(x,hD_x) \\ &\quad=\frac{i}{h}\Big[h^2D_{x_n}^2\psi_h^w(hD_{x_n}),\ \chi(x_n)\,hD_{x_n}\,\chi_{in}^w(\frac{R(x',0,\xi')-1}{h^\delta})\Big] \\ &\qquad+\frac{i}{h}\Big[R(x',x_n,hD_{x'}),\ \chi(x_n)\,hD_{x_n}\,\chi_{in}^w(\frac{R(x',0,\xi')-1}{h^\delta})\Big] \end{aligned} \tag{3.8}$$

$$\quad+\mathcal{O}_\delta(h^\infty)_{L^2\to L^2}=:b_1^w(x,hD_x)+b_2^w(x,hD_x). \tag{3.9}$$

Clearly,

$$b_1^w(x,hD_x)=\frac{i}{h}\Big[h^2D_{x_n}^2\psi_h^w(hD_{x_n}),\ \chi(x_n)\,hD_{x_n}\,\chi_{in}^w(\frac{R(x',0,\xi')-1}{h^\delta})\Big]\in Op_h^w(S^0_{\Sigma_H,\delta})$$

with

$$\sigma(b_1)(x,\xi;h)=2\xi_n^2\psi(\xi_n)\chi'(x_n)\chi_{in}^w(\frac{R(x',0,\xi')-1}{h^\delta}).$$

As for the second term,

$$\frac{i}{h}\Big[R(x_n,x',hD_{x'}),\chi(x_n)hD_{x_n}\chi_{in}^w(\frac{R(x',0,\xi')-1}{h^\delta})\Big]\in Op_h^w(S^\delta_{\Sigma_H,\delta}).$$

Thus, $b^w(x,hD_x) \in Op_h^w(S^{\delta}_{\Sigma_H,\delta})$ and by L^2-boundedness (see (2.10)),

$$b^w(x,hD_x) = b_2^w(x,hD_x) + \mathcal{O}_\delta(1)_{L^2 \to L^2}. \tag{3.10}$$

Finally, we simply note that exactly the same argument holds for χ_{tan} replacing χ_{in} in the above.

$\square$

4. ESTIMATING INTERIOR TERMS

4.1. **Dirichlet data.** From Theorem 1.1 we have with $b_{in} \in Op_h(S^{\delta}_{\Sigma_H,\delta})$,

$$\begin{aligned} &\Big\langle (1+h^2\Delta_H)(\chi_{in})^w_{h,\delta}\varphi_h^H, \varphi_h^H \Big\rangle_{L^2(H)} + \Big\langle (\chi_{in})^w_{h,\delta}\varphi_h^{H,\nu}, \varphi_h^{H,\nu} \Big\rangle_{L^2(H)} \\ &= \Big\langle b^w_{in}(x,hD_x)\varphi_h, \varphi_h \Big\rangle_{L^2(M_-)} + \mathcal{O}(h^\infty) = \mathcal{O}_\delta(h^{-\delta}), \end{aligned} \tag{4.1}$$

where the last line in (4.1) follows by L^2-boundedness of $h^\delta b^w_{in}(x,hD_x) \in Op_h^w(S^0_{\Sigma_H,\delta})$. Moreover since $1 - R(x', x_n = 0, \xi') \gtrapprox h^\delta$ for $(x',\xi') \in \text{supp}(\chi_{in})_\delta$, it follows by the sharp Gårding inequality that

$$\Big\langle (1+h^2\Delta_H)(\chi_{in})^w_{h,\delta}\varphi_h^H, \varphi_h^H \Big\rangle_{L^2(H)} \gtrapprox h^\delta \Big\langle (\chi_{in})^w_{h,\delta}\varphi_h^H, \varphi_h^H \Big\rangle_{L^2(H)}. \tag{4.2}$$

Since the normal derivative term on the LHS of (4.1) is non-negative, one gets the interior Dirichlet bound

$$\Big\langle (\chi_{in})^w_{h,\delta}\varphi_h^H, \varphi_h^H \Big\rangle_{L^2(H)} = O_\delta(h^{-2\delta}). \tag{4.3}$$

4.2. **Neumann data.** In view of the positivity of (4.2), it follows from the $O(h^{-\delta})$-bound for the RHS of (4.1) that for the Neumann data one has the better bound

$$\Big\langle (\chi_{in})^w_{h,\delta}\varphi_h^{H,\nu}, \varphi_h^{H,\nu} \Big\rangle_{L^2(H)} = O_\delta(h^{-\delta}). \tag{4.4}$$

5. ESTIMATING THE EXTERIOR TERMS

5.1. **Exterior mass estimates on $\mathcal{O}(1)$-scales.** A key issue is mass concentration of eigenfunctions: Let $\varphi_{\lambda_j}; j = 1,2,...$ be an L^2 orthonormal basis of Laplace eigenfuncgtions on (M,g) and $H \subset M$ a hypersurface. To begin we first discuss mass concentration of φ_h^H on B^*H on scales ~ 1 and then indicate the extension to small-scales $\sim h^\delta; \delta \in [0,1)$ using 2-microlocal calculus in subsection 5.2.

Let $h \in \{\lambda_j^{-1}\}; j = 1,2,....$ Given $0 < \epsilon_0 < 1$ an arbitrary small number, let $\chi(x,\xi) \in C_0^\infty(T^*M)$ be equal to one on the annulus $A(\epsilon_0) = \{(x,\xi); (1-\epsilon_0) < |\xi|_g < (1+\epsilon)$ and with supp $\chi \subset A(2\epsilon_0)$. Let $\tilde{\chi} \in C_0^\infty$ be another cutoff equal to one on $A(2\epsilon_0)$ and with supp $\tilde{\chi} \subset A(4\epsilon_0)$. Consider the eigenfunction equation

$$(-h^2\Delta_g - 1)\varphi_h = 0.$$

Then, $P(h) := -h^2\Delta_g - 1$ is h elliptic for $(x,\xi) \in T^*M - A(\epsilon)$. So, one can construct an h-microlocal parametrix with $Q(h) \in Op_h(S^0_{0,0})$ so that

$$(1-\tilde{\chi}(h))Q(h)P(h)(1-\chi(h))\varphi_h = (1-\tilde{\chi}(h))\varphi_h + \mathcal{O}(h^\infty).$$

Since $P(h)\varphi_h = 0$ and $\sigma([P(h),(1-\chi(h))](x,\xi) = 0$ for $(x,\xi) \in$supp $(1-\tilde{\chi}(h))$, one gets the well-known concentration estimate

$$\|(1-\tilde{\chi}(h))\varphi_h\|_{L^2} = \mathcal{O}(h^\infty). \tag{5.1}$$

A similar argument with the derivatives $\partial_x^\alpha \varphi_h(x)$ combined with Sobolev embedding implies

$$\|(1-\tilde{\chi}(h))\varphi_h\|_{C^k} = \mathcal{O}(h^\infty), \tag{5.2}$$

and as a consequence

$$WF_h(\varphi_h) \subset S^*M.$$

Proposition 5.1. *Let* $H \subset M$ *be a hypersurface and* $\varphi_h^H := \varphi_h|_H = \gamma_H \varphi_h$ *and* $\varphi_h^{H,\nu} = \gamma_H(hD_\nu \varphi_h)$ *as usual. Then,*

$$WF_h(\varphi_h^H) \subset \overline{B^*H}$$

and

$$WF_h(\varphi_h^{H,\nu}) \subset \overline{B^*H}.$$

Proof. Let $(x', x_n) \in \mathbb{R}^{n-1} \times (-\epsilon_0, \epsilon_0)$ be Fermi coordinates in an ϵ_0 collar neighbourhood near H with $H = \{x_n = 0\}$. In these coordinates,

$$-h^2\Delta_g = h^2 D_{x_n}^2 - h^2 \Delta_H + \mathcal{O}(x_n)|hD_{x'}|^2 + \mathcal{O}(h^2(|D_x'| + |D_{x_n}|)). \tag{5.3}$$

Let $\zeta(x', x_n, \xi') \in C_0^\infty(\mathbb{R}^{n-1} \times (-\epsilon_0, \epsilon_0) \times \mathbb{R}^{n-1})$ be equal to 1 when $\xi' \in B(1+\epsilon_0)$ and vanishing outside $B(1+2\epsilon_0)$. As above, we denote by $\tilde{\zeta}$ a cutoff with bigger support, equal to 1 in $B(1+2\epsilon_0)$ and vanishing outside $B(1+4\epsilon_0)$. Since $[\gamma_H, \Delta_H] = 0$, for $j = 0, 1$ we have

$$(-h^2\Delta_H - 1)(1 - \zeta(x', x_n = 0, hD_{x'}))\gamma_H((hD_n)^j \varphi_h) \tag{5.4}$$

$$\begin{aligned} &= \gamma_H\, [-h^2 D_{x_n}^2 + \mathcal{O}(x_n)|hD_{x'}|^2 \\ &\quad + \mathcal{O}(h^2(|D_x'| + |D_{x_n}|))\,](1 - \zeta(x', x_n, hD_{x'}))(hD_n)^j \varphi_h \\ &\quad + \gamma_H(-h^2\Delta_g - 1)(1 - \zeta(x', x_n, hD_{x'}))(hD_n)^j \varphi_h. \end{aligned} \tag{5.5}$$

Since $|\xi'|^2 \geq 1 + 2\epsilon_0$ on supp $(1 - \tilde{\zeta}(x', x_n, \xi'))$, obviously $\xi_n^2 + |\xi'|^2 \geq 1 + 2\epsilon_0$ also holds on supp $1 - \tilde{\zeta}$. But then, by (5.2) (and since an application of hD_n does not increase the wavefront set) it follows that both terms on the RHS of (5.4) are $\mathcal{O}(h^\infty)$. As for the LHS, it then follows that

$$(1 - \tilde{\zeta}(x', x_n = 0, hD_{x'})) \cdot (-h^2\Delta_H - 1) \cdot (1 - \zeta(x', x_n = 0, hD_{x'}))\varphi_h^H = \mathcal{O}(h^\infty).$$

Then, since $(1-\zeta)(h^2\Delta_H - 1)$ is h-elliptic on supp $1 - \tilde{\zeta}(x_n = 0, x', \xi')$, by the same kind of parametrix construction used to prove (5.1), it follows that

$$\|(1 - \tilde{\zeta}(x', x_n = 0, hD_{x'}))\gamma_H((hD_n)^j \varphi_h)\|_{L^2} = \mathcal{O}(h^\infty).$$

By a Sobolev argument, the same bound holds for C^k-norms. □

5.2. **Mass estimates on scales $\mathcal{O}(h^\delta)$.** We only need modify the argument in subsection 5.1 by choosing appropriate pseudodifferential cutoffs in $Op_h^w(S^0_{\Sigma_H,\delta})(T^*M)$ and then apply an analogous 2-microlocal parametrix construction. More precisely, we define

$$\zeta(x', x_n, hD_{x'}) := \chi(\epsilon_0^{-1} h^{-\delta} x_n) \cdot (\chi_{out})^w_{h,\delta}(x', hD_{x'}).$$

Then, by choosing $\epsilon_0 > 0$ small, for $(x', x_n, \xi') \in \text{supp}(\zeta)$ we have that

$$|\xi|_g^2 = \xi_n^2 + R(x', x_n, \xi') = \xi_n^2 + R(x', x_n = 0, \xi') + \mathcal{O}_{\epsilon_0}(h^\delta)$$

$$\geq \xi_n^2 + 1 + \frac{1}{2}h^\delta \geq 1 + \frac{1}{2}h^\delta$$

Using the fact that operators in $Op_h^w(S^*_{\Sigma_H,\delta})$ constitute an algebra (see Proposition 2.2), the argument then proceeds as in the proof of Proposition 5.1 and one gets that

$$\max\left(\left\langle(\chi_{out})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle,\left\langle(\chi_{out})^w_{h,\delta}\varphi_h^{H,},\varphi_h^{H}\right\rangle\right)=\mathcal{O}_\delta(h^\infty). \tag{5.6}$$

6. Proof of Theorem 1

Proof. Since

$$\begin{aligned}\|\varphi_h^{H,\nu}\|^2_{L^2(H)}&=\left\langle(\chi_{in})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle_{L^2(H)}+\left\langle(\chi_{tan})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle_{L^2(H)}\\&\quad+\left\langle(\chi_{out})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle_{L^2(H)},\end{aligned}$$

in view of the bounds in (4.4) and (5.6), it follows that

$$\|\varphi_h^{H,\nu}\|^2_{L^2(H)}=\left\langle(\chi_{tan})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle_{L^2(H)}+O(h^{-\delta}). \tag{6.1}$$

To estimate the tangential term on the RHS of (6.1) we note the difficulty is in estimating a term of the form

$$\left\langle(1+h^2\Delta_H)(\chi_{tan})^w_{h,\delta}\varphi_h^{H},\varphi_h^{H}\right\rangle_{L^2(H)},$$

which is no longer positive since $1+h^2\Delta_H$ is small on the microsupport of $(\chi_{tan})^w_{h,\delta}$. To estimate this tangential term, we use Theorem 1.1 yet again to get that

$$\begin{aligned}&\left\langle(1+h^2\Delta_H)(\chi_{tan})^w_{h,\delta}\varphi_h^{H},\varphi_h^{H}\right\rangle_{L^2(H)}+\left\langle(\chi_{tan})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle_{L^2(H)}\\&=\left\langle b^w_{tan}(x,hD_x)\varphi_h,\varphi_h\right\rangle_{L^2(M_-)}+\mathcal{O}(h^\infty)=\mathcal{O}_\delta(\hbar^{-\delta}),\end{aligned} \tag{6.2}$$

Since $|R(x',x_n=0,\xi')-1|\lessapprox h^\delta$ when $(x',\xi')\in\text{supp}\,(\chi_{tan})_{h,\delta}$, by L^2-boundedness,

$$\left\langle(1+h^2\Delta_H)(\chi_{tan})^w_{h,\delta}\varphi_h^{H},\varphi_h^{H}\right\rangle_{L^2(H)}=\mathcal{O}_\delta(h^\delta)\|\varphi_h^H\|^2_{L^2(H)}. \tag{6.3}$$

It follows by the universal bounds of Burq-Gérard-Tzvetkov [BGT] that $\|\varphi_h^H\|_{L^2(H)}=O(h^{-1/4})$. Thus, from (6.3) and (6.2),

$$\left\langle(\chi_{tan})^w_{h,\delta}\varphi_h^{H,\nu},\varphi_h^{H,\nu}\right\rangle_{L^2(H)}=\mathcal{O}_\delta(h^{\delta-\frac{1}{2}}+h^{-\delta}). \tag{6.4}$$

So, by (6.1) we get that

$$\|\varphi_h^{H,\nu}\|^2_{L^2(H)}=O_\delta(h^{\delta-\frac{1}{2}}+h^{-\delta}). \tag{6.5}$$

Choosing $\delta=\frac{1}{4}$ to optimize (6.5) finishes the proof of Theorem 1. $\square$

References

[Bur] Burq, N., Quantum ergodicity of boundary values of eigenfunctions: a control theory approach. *Canad. Math. Bull.* **48** (2005), no. 1, 3–15.

[BGT] Burq, N., Gérard, P., and Tzvetkov, N. Restrictions of the Laplace-Beltrami eigenfunctions to submanifolds. *Duke Math. J.* **138** (2007), no. 3, 445-486.

[CTZ] Christianson, H., Toth, J.A., and Zelditch, S. Quantum Ergodic Restriction for Cauchy Data: Interior QUE and Restricted QUE. *submitted*, (arXiv:1205.0286).

[CTZ2] Christianson, H., Toth, J.A., and Zelditch, S. δ-Quantum Ergodicity and Restrictions of Eigenfunctions *in preparation*.

[HT] Hassell, A. and Tao, T. Upper and lower bounds for normal derivatives of Dirichlet eigenfunctions. *Math. Res. Lett.* **9** (2002), no. 2–3, 289-305.

[SjZw1] Sjöstrand, J. and Zworski, . Asymptotic distribution of resonances for convex obstacles. *Acta Math.* **183** (1999), no. 2, 191–253.

[SjZw2] Sjöstrand, J. and Zworski, M. Fractal upper bounds on the density of semiclassical resonances. *Duke Math. J.* **137** (2007), no. 3, 381–459.

[TZ2] Toth, J. and Zelditch, S. Quantum ergodic restriction thoerems, manifolds without boundary. GAFA (2013)(to appear), (arXiv:1104.4531).

[Zw] Zworski, M. Semiclassical Analysis. Graduate Studies in Mathematics, 138. *American Mathematical Society, Providence, RI*, 2012.

E-mail address: `hans@math.unc.edu`

Department of Mathematics, UNC-Chapel Hill, CB#3250 Phillips Hall, Chapel Hill, NC 27599

Department of Mathematics, McGill University, Montreal, CANADA
E-mail address: `jtoth@math.mcgill.ca`